Dept of Mathematics and Statistics
King Fahd University of Petroleum & Minerals
AS481: Actuarial Contingencies II
Dr. Ridwan A. Sanusi
Major Exam I, Term 261
Monday, Sept. 14, 2026
1:00 PM – 2:50 PM

Name……. ID#:_____ Serial #:___

Instructions.

1. Any student caught with mobile phones on during the exam will be considered under the cheating rules of the University.
2. If you need to leave the room, please do so before the exam starts. Nobody will be allowed to leave the room once the exam starts.
3. Only materials provided by the instructor can be present on the table during the exam.
4. Do not spend too much time on any one question. If a question seems too difficult, leave it and go on.
5. Use the blank portions of each page for your work. Extra blank pages can be provided if necessary. If you use an extra page, indicate clearly what problem you are working on.
6. ***Only answers supported by work will be considered. Unsupported guesses will not be graded.***
7. While every attempt is made to avoid defective questions, sometimes they do occur. In the rare event that you believe a question is defective, the instructor cannot give you any guidance beyond these instructions.
8. Mobile calculators, I-pad, or communicable devices are disallowed. Use regular scientific calculators, financial calculators, or SOA approved calculators only. ***Write important steps to arrive at the solution of the exam problems.***

The test is 110 minutes, GOOD LUCK, and you may begin now!

Question	**Total Marks**	**Marks Obtained**	**Comments**
1-10	5		
11-15	5		
PART B			
1	2.5		
2	2.5		
Total	15		

Extra blank page

PART A

1. In a discrete-time Markov chain, which of the following best describes the memoryless property?

A) The process can only move to adjacent states in the next time step
B) The probability of moving to the next state depends on the entire history of past states
C) The probability of moving to the next state depends only on the current state
D) The process must eventually return to its starting state
E) The transition probabilities must be the same for all states.

2. Markov chain has transition probability matrix $\boldsymbol{P}$, with states 1, 2, ….. If State 2 is an absorbing state, which of the following must be true?

A) The second column of $\boldsymbol{P}$ has all entries equal to 0
B) The second row of $\boldsymbol{P}$ has $p^{22} = 1$ and all other entries in that row equal to 0
C) The second row of $\boldsymbol{P}$ has all entries equal to 0
D) The second column of $\boldsymbol{P}$ has $p^{22} = 1$ and all other entries in that column equal to 0
E) State 2 cannot communicate with any other state, but can be left with probability 1.

3. A non-homogeneous discrete-time Markov chain is characterized by:

A) The transition probability matrix $\boldsymbol{P}$ is the same for every time interval
B) The state space changes over time
C) The transition probability matrix may differ for each time interval, denoted $\boldsymbol{P}^{(k)}$
D) The process has no absorbing states
E) The number of states increases with time.

4. A two-state homogeneous Markov chain has transition probability matrix:

$$\boldsymbol{P} = \begin{bmatrix} 0.3 & 0.7 \\ 0.4 & 0.6 \end{bmatrix}$$

If the process is in State 0 at time 0, what is the probability that it is in State 0 at time 2?

A) 0.09
B) 0.21
C) 0.28
D) 0.37
E) 0.49.

5. In the two-state continuous-time Markov survival model (State 0 = Alive, State 1 = Dead), which of the following statements about the probability ${}_tp_x^{00}$ is correct?

A) ${}_tp_x^{00}$ represents the probability of leaving State 0 by time t
B) ${}_tp_x^{00}$ represents the probability of being in State 0 at time t
C) ${}_tp_x^{00}$ represents the probability of being in State 1 at time t
D) ${}_tp_x^{00}$ represents the probability of transitioning from State 0 to State 1 within time t
E) ${}_tp_x^{00}$ is always equal to ${}_tq_x$.

6. In the previous question, the Kolmogorov forward equation for ${}_tp_x^{00}$ is:

$$\frac{d}{dt}\ {}_tp_x^{00} = -\ {}_tp_x^{00} \cdot \mu_{x+t}^{01}$$

Why does this equation contain no positive term representing probability flowing back into State 0?

A) Because the force of mortality μ_{x+t}^{01} is constant
B) Because State 1 is absorbing, so the force of transition from State 1 to State 0 is zero
C) Because the process is homogeneous
D) Because ${}_tp_x^{00}$ is always equal to 1
E) Because the model has only one state.

7. In a multiple-decrement model, which of the following correctly distinguishes the dependent probability $q_x^{(j)}$ from the independent probability $q_x'^{(j)}$?

A) $q_x^{(j)}$ is always greater than $q_x'^{(j)}$
B) $q_x^{(j)}$ is the probability of decrement from Cause j in the presence of other causes, while $q_x'^{(j)}$ is the probability if Cause j were the only decrement operating
C) $q_x'^{(j)}$ is the probability of decrement from Cause j in the presence of other causes, while $q_x^{(j)}$ is the probability if Cause j were the only decrement operating
D) $q_x^{(j)}$ and $q_x'^{(j)}$ are always equal
E) $q_x^{(j)}$ applies only to continuous models, while $q_x'^{(j)}$ applies only to discrete models.

8. Which of the following is an example of a competing risks scenario in actuarial science?

A) A pension plan where a member may retire, die, or leave for another job, but only one of these events will occur
B) A policyholder simultaneously holding multiple insurance policies
C) A medical plan that provides coverage for two different treatments
D) All of the above.
E) None of the above.

9. In a discrete-time Markov chain with state space $\{1,2,3,\ldots,m\}$, let

$\pi_n = (\pi_{1n}, \pi_{2n}, \ldots, \pi_{mn})$. Which of the statements is correct?

A. π_n is the transition matrix at time n, and $\pi_{in}$is the probability of moving from state i to state n.

B. π_n is the distribution of *first passage times*, and π_{in} is the probability that the process first enters state $i$at time n.

C. π_n is the vector of state-specific expected rewards at time n, and π_{in} gives the expected reward of being in state $i$at time n.

D. π_n is the state probability vector at time n, and π_{in} is the probability that the chain is in state i at time n.

E. π_n is the vector of stationary probabilities, and $\pi_{in}$is the long-run proportion of transitions into state $i$over all time.

10. A certain animal species can be classified as thriving (State 0), endangered (State 1), or extinct (State 2). Movement among states is governed by a non-homogeneous Markov process defined by the following transition probability matrices:

$$\mathbf{P}^{(0)} = \begin{bmatrix} .85 & .15 & 0 \\ 0 & .70 & .30 \\ 0 & 0 & 1 \end{bmatrix}, \qquad \mathbf{P}^{(1)} = \begin{bmatrix} .90 & .10 & 0 \\ .10 & .70 & .20 \\ 0 & 0 & 1 \end{bmatrix},$$

$$\mathbf{P}^{(2)} = \begin{bmatrix} .95 & .05 & 0 \\ .20 & .70 & .10 \\ 0 & 0 & 1 \end{bmatrix}, \qquad \mathbf{P}^{(k)} = \begin{bmatrix} .95 & .05 & 0 \\ .50 & .50 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

where $k = 3,4,5,\cdots$.

If the species is thriving at time n, is it possible for it to be extinct at time n +1?

A. Yes
B. No
C. Not enough information to decide
D. It's confusing
E. None of the above.

Show workings for Questions 11 – 15

11. A discrete-time Markov process has only two states, denoted as State 0 and State 1. A person is age x at time 0, and is located in State 0. We are given the age-specific one-year transition probability values

$$p_{x+k}^{00} = .70 + \frac{.10}{k+1}$$

and

$$p_{x+k}^{11} = .60 + \frac{.20}{k+1}.$$

Find the value of Pr [X3 = 0 | X1 = 0].

A. 0.6333
B. 0.2500
C. 0.3333
D. 0.3145
E. 0.7500

12.

If $\mu_{x+t}^{(1)} = .10$ and $\mu_{x+t}^{(2)} = .20$ for all t, find ${}_{\infty}q_x^{(1)}$

A. 0.3333
B. 0.6667
C. 0.2000
D. 0.8000

13. Given the following probabilities of decrement. Assume an initial group (called the radix of the table) of size 1000. Calculate the probability that a person in the group at age 46 will leave between ages 48 and 49.

x	$q_x^{(1)}$	$q_x^{(2)}$
45	.011	.100
46	.012	.100
47	.013	.100
48	.014	.100
49	.015	.100
50	.016	.100

A. 0.0110
B. 0.6204
C. 0.6979
D. 0.0201
E. 0.0898

14. In Question 13, calculate ${}_{2|2}q_{45}^{(1)}$.
A. 0.0110
B. 0.6204
C. 0.6979
D. 0.0201
E. 0.0898

15. In Question 13, calculate $Pr(K_x = 2 \cap J_x = 1)$.
A. 0.0110
B. 0.6204
C. 0.6979
D. 0.0201
E. 0.0898

PART B

1.

Students can leave a certain three-year school only for reasons of failure (Decrement 1) or voluntary withdrawal (Decrement 2), where each decrement is uniformly distributed over $(x, x+1)$ in its associated single-decrement table. The following values are given:

x	$q'^{(1)}_x$	$q'^{(2)}_x$	$q^{(1)}_x$	$q^{(2)}_x$
0	.10	.25	–	–
1	.20	.20	–	–
2	.20	.10	–	–

A. Calculate the six missing probability values for the table.

B. Find the marginal probabilities $p_{K_0}(1)$, $p_{K_0}(2)$, $p_{J_0}(1)$ and $p_{J_0}(2)$.

2.

The following double-decrement table gives probability values for a student at the beginning of each year in a three-year Actuarial Science Graduate School. Some of the entries in the table have been obliterated by tear stains.

Academic Year	Probability of Academic Failure	Probability of Voluntary Withdrawal	Probability of Completing the Year
1	.40	.20	–
2	–	.30	–
3	–	–	.60

It is known that ten times as many students complete Year 2 as fail during Year 3, and the number of students who fail during Year 2 is 40% of the number who complete Year 2. Find the probability that a new student entering the school will voluntarily withdraw before graduation.

Chapter 3 MQR Markov Chain review

$p^{ij} = P\left(X_{n+1} = j | X_n = i\right) \quad \sum_{j=0}^{m} p^{ij} = 1 \quad \mathbf{P} = \begin{bmatrix} p_{00} & p_{01} & \cdots & p_{0m} \\ p_{10} & p_{11} & \cdots & p_{1m} \\ \vdots & \vdots & \ddots & \vdots \\ p_{m0} & p_{m1} & \cdots & p_{mm} \end{bmatrix}$

$\boldsymbol{\pi}_n = (\pi_{0n}, \pi_{1n}, \cdots, \pi_{mn}) \quad \sum_{i=0}^{m} \pi_{in} = 1 \quad \boldsymbol{\pi}_{n+r} = \begin{cases} \boldsymbol{\pi}_n \cdot \mathbf{P} \cdot \mathbf{P} \cdot \cdots \cdot \mathbf{P} = \boldsymbol{\pi}_n \cdot \mathbf{P}^r & \text{homogeneous} \\ \boldsymbol{\pi}_n \cdot \mathbf{P}^{(0)} \cdot \mathbf{P}^{(1)} \cdot \cdots \cdot \mathbf{P}^{(r-1)} & \text{non-homogeneous} \end{cases}$

${}_rp_n^{ij} = P\left(X_{n+r} = j | X_n = i\right) \qquad {}_rp_n^{\overline{ii}} \leq {}_rp_n^{ii}$

${}_rp_n^{\overline{ii}} = {}_rp_n^{ii}$ if **cannot** reenter state i once left i

μ_{x+s}^{ij} =force of transition. $\quad \mu_{x+s}^{i} = \sum_{j=1}^{i-1} \mu_{x+s}^{ij} + \sum_{j=i+1}^{m} \mu_{x+s}^{ij}$ =force of transitioning **out**.

$$\frac{d}{dr}({}_rp_x^{ij}) = \sum_{k \neq j} \left({}_rp_x^{ik} \mu_{x+r}^{kj} - {}_rp_x^{ij} \cdot \mu_{x+r}^{jk} \right) = \sum_{k \neq j} \left({}_rp_x^{ik} \mu_{x+r}^{kj} \right) - {}_rp_{ij}^{(t)} \cdot \mu_{x+r}^{i}$$

Chapter 13 MQR Multiple Decrement Models: Theory

OBJECTIVES: 1.To understand the concept of a multiple decrement table

2.To understand the force of decrement

3.To construct a multiple decrement model using associated single decrements and to apply various assumptions to calculate rates for discrete jumps.

13.1 Discrete Multiple Decrement Models

$q_x^{(\tau)} = q_x^{(1)} + q_x^{(2)} + \ldots + q_x^{(m)} = \sum_{j=1}^{m} q_x^{(j)} \quad (13.1) \qquad p_x^{(\tau)} = 1 - q_x^{(\tau)} \quad (13.2)$

$d_x^{(\tau)} = \sum_{j=1}^{m} d_x^{(j)} = \ell_x^{(\tau)} \cdot q_x^{(\tau)} \quad (13.3 \text{ \& } 13.7b) \qquad {}_np_x^{(\tau)} = 1 - {}_nq_x^{(\tau)} \quad (13.7e)$

${}_nd_x^{(j)} = \sum_{t=0}^{n-1} d_{x+t}^{(j)} = \ell_x^{(\tau)} \cdot {}_nq_x^{(j)} \quad (13.4 \text{ \& } 13.7c) \qquad \ell_x^{(\tau)} = \sum_{j=1}^{m} \ell_x^{(j)} \quad (13.6)$

$d_x^{(j)} = \ell_x^{(\tau)} \cdot q_x^{(j)} \quad (13.7a) \qquad \ell_{x+n}^{(\tau)} = \ell_x^{(\tau)} \cdot {}_np_x^{(\tau)} \quad (13.7f)$

${}_nd_x^{(\tau)} = \sum_{j=1}^{m} {}_nd_x^{(j)} = \ell_x^{(\tau)} \cdot {}_nq_x^{(\tau)} \quad (13.5a \text{ \& } 13.7d) \qquad {}_nq_x^{(\tau)} = {}_nd_x^{(\tau)} / \ell_x^{(\tau)} = \sum_{j=1}^{m} {}_nq_x^{(j)} \quad (13.5b)$

13.1.2 Random Variable Analysis

The *joint probability function* of K_x and J_x is $\Pr\left(K_x = k \cap J_x = j\right) = {}_k|\, q_x^{(j)} = \frac{d_{x+k}^{(j)}}{\ell_x^{(\tau)}} \quad (13.8)$

The *marginal probability functions* are

i) $\Pr\left(K_x = k\right) = \sum_{j=1}^{m} \Pr\left(K_x = k \cap J_x = j\right) = {}_k|\, q_x^{(\tau)} = \frac{d_{x+k-1}^{(1)} + \ldots + d_{x+k}^{(m)}}{\ell_x^{(\tau)}} = \sum_{j=1}^{m} \frac{d_{x+k}^{(j)}}{\ell_x^{(\tau)}} \quad (13.9)$

ii) $\Pr\left(J_x = j\right) = \sum_{k=1}^{\infty} \Pr\left(K_x = k \cap J_x = j\right) = \sum_{k=1}^{\infty} \frac{d_{x+k}^{(j)}}{\ell_x^{(\tau)}} \quad (13.10)$

13.2 Theory of Competing Risks ${}_nq_x'^{(j)} \geq {}_nq_x^{(j)} \quad (13.11)$

13.3 Continuous Multiple Decrement Models

$\mu_{x+t}^{(j)} = \frac{-\frac{d}{dt}\, {}_tp_x'^{(j)}}{{}_tp_x'^{(j)}} \quad (13.12a) \qquad \mu_{x+t}^{(\tau)} = \frac{-\frac{d}{dt}\, {}_tp_x^{(\tau)}}{{}_tp_x^{(\tau)}} \quad (13.13a)$

${}_tq_x'^{(j)} = \int_0^t {}_sp_x'^{(j)} \mu_{x+s}^{(j)}\, ds = 1 - {}_tp_x'^{(j)} \qquad {}_tp_x'^{(j)} = \exp\left(-\int_0^t \mu_{x+s}^{(j)}\, ds \right) \quad (13.12b)$

$f_{x^{(j)}}(t) = {}_tp_x'^{(j)} \cdot \mu_{x+t}^{(j)} \quad (13.14) \qquad {}_tp_x^{(\tau)} = \exp\left(-\int_0^t \mu_{x+s}^{(\tau)}\, ds \right) \quad (13.13b)$

$F_{x^{(j)}}(t) = \Pr\left[T_x^{(j)} \leq t\right] = \int_0^t f_{x^{(j)}}(s)\, ds = \int_0^t {}_sp_x'^{(j)} \cdot \mu_{x+s}^{(j)}\, ds \quad (13.15)$

$$\begin{aligned} \mu_{x+t}^{(\tau)} &= \frac{-\frac{d}{dt}\, {}_tp_x^{(\tau)}}{{}_tp_x^{(\tau)}} = -\frac{d}{dt} \ln {}_tp_x^{(\tau)} = -\frac{d}{dt} \ln \left[{}_tp_x'^{(1)} \cdot {}_tp_x'^{(2)} \cdot \ldots \cdot {}_tp_x'^{(m)} \right] \\ &= \left(-\frac{d}{dt} \ln {}_tp_x'^{(1)} \right) + \left(-\frac{d}{dt} \ln {}_tp_x'^{(2)} \right) + \ldots + \left(-\frac{d}{dt} \ln {}_tp_x'^{(m)} \right) = \mu_{x+t}^{(1)} + \mu_{x+t}^{(2)} + \ldots + \mu_{x+t}^{(m)} = \sum_{j=1}^{m} \mu_{x+t}^{(j)} \quad (13.17) \end{aligned}$$

Fundamental Relation Between Primed and Unprimed Rates: ${}_tp_x^{(\tau)} = \exp\left(-\sum_{j=1}^{m}\int_0^t \mu_{x+s}^{(j)}\,ds\right) = \prod_{j=1}^{m} {}_tp_x^{\prime(j)}$ (13.16)

${}_tq_x^{(j)} = \int_0^t f_{T,J}(s,j)\,ds = \int_0^t {}_sp_x^{(\tau)}\cdot\mu_{x+s}^{(j)}\,ds$ (13.18) ${}_tq_x^{(\tau)} = \int_0^t {}_sp_x^{(\tau)}\cdot\mu_{x+s}^{(\tau)}\,ds$ (13.20)

$\frac{d}{dt}\,{}_tq_x^{(j)} = \frac{d}{dt}\int_0^t {}_sp_x^{(\tau)}\cdot\mu_{x+s}^{(j)}\,ds = {}_tp_x^{(\tau)}\cdot\mu_{x+s}^{(j)} \;\rightarrow\; \mu_{x+t}^{(j)} = \frac{\frac{d}{dt}\,{}_tq_x^{(j)}}{{}_tp_x^{(\tau)}}$ (13.19)

Joint Distribution of T_x and J_x $\Pr(t < T_x \le t+dt \text{ and } J_x = j) \approx {}_tp_x^{(\tau)}\mu_{x+t}^{(j)}dt$, ${}_tq_x^{(j)} = \int_0^t {}_sp_x^{(\tau)}\,\mu_{x+s}^{(j)}\,ds$.

13.4.1 Uniform Distribution of Decrements in the Multiple Decrement Context

${}_tq_x^{(j)} = t\cdot q_x^{(j)}$ (13.21) $q_x^{(j)} = {}_tp_x^{(\tau)}\cdot\mu_{x+t}^{(j)}$ (13.22)

${}_tq_x^{(\tau)} = t\cdot q_x^{(\tau)}$ (13.23) ${}_tp_x^{(\tau)} = 1 - t\cdot q_x^{(\tau)}$ (13.24)

$\mu_{x+t}^{(j)} = \frac{q_x^{(j)}}{{}_tp_x^{(\tau)}} = \frac{q_x^{(j)}}{1-t\cdot q_x^{(\tau)}}$ (13.25) ${}_tp_x^{\prime(j)} = \exp\left[\frac{q_x^{(j)}}{q_x^{(\tau)}}\cdot\ln\left(1-t\cdot q_x^{(\tau)}\right)\right] = \left(1-t\cdot q_x^{(\tau)}\right)^{q_x^{(j)}/q_x^{(\tau)}}$ (13.26)

13.4.2 Uniform Distribution in the Associated Single- Decrement Tables

case ${}_tq_x^{\prime(j)} = t\cdot q_x^{\prime(j)}$ (13.27) ${}_tp_x^{\prime(j)}\cdot\mu_{x+t}^{(j)} = q_x^{\prime(j)}$ (13.28)

Double decrement	$q_x^{(1)} = \int_0^1\left(1-t\cdot q_x^{\prime(2)}\right)\cdot q_x^{\prime(1)}dt = q_x^{\prime(1)}\left(1-\frac{1}{2}\cdot q_x^{\prime(2)}\right)$ (13.29a)	$q_x^{(2)} = q_x^{\prime(2)}\left(1-\frac{1}{2}\cdot q_x^{\prime(1)}\right)$ (13.29b)
Triple Decrement	$q_x^{(1)} = q_x^{\prime(1)}\left[1-\frac{1}{2}\left(q_x^{\prime(2)}+q_x^{\prime(3)}\right)+\frac{1}{3}\left(q_x^{\prime(2)}\cdot q_x^{\prime(3)}\right)\right]$ (13.30)	

13.4.3 Constant Force of Decrement

$\frac{{}_tq_x^{(j)}}{{}_tq_x^{(\tau)}} = \frac{\mu_{x+t}^{(j)}}{\mu_{x+t}^{(\tau)}}$

Miscellaneous Results (From ACTEX MLC manual)

1. Assumptions on the single decrement table.

Backing out the Unprimed Rates from Primed Rates

${}_sq_x^{(i)} = \int_0^s {}_tp_x^{(\tau)}\mu_{x+t}^{(i)}\,dt = \int_0^s\left[\prod_{j=1,\,j\neq i}^{m} {}_tp_x^{\prime(j)}\right]{}_tp_x^{\prime(i)}\mu_{x+t}^{(i)}\,dt$

$$\frac{d}{dr}({}_rp_x^{ij}) = \sum_{k\neq j}\left({}_rp_x^{ik}\mu_{x+r}^{kj} - {}_rp_x^{ij}\cdot\mu_{x+r}^{jk}\right) = \sum_{k\neq j}\left({}_rp_x^{ik}\mu_{x+r}^{kj}\right) - {}_rp_{ij}^{(t)}\cdot\mu_{x+r}^{i}$$

2. **Constant Force Assumption for Multiple Decrements** For any $t\in[0,1]$ and integer-valued x,

(i) ${}_tp_x^{(\tau)} = \left[p_x^{(\tau)}\right]^t$ (survival probability for **fractional** ages)

(ii) *Ratio* Property : $\frac{{}_tq_x^{(i)}}{{}_tq_x^{(\tau)}} = \frac{\mu_{x+s}^{(i)}}{\mu_{x+s}^{(\tau)}}$ for any $s\in[0,1]$ (To get unprimed rates from (i))

(iii) *Partition* Property : ${}_tp_x^{t(i)} = \left[{}_tp_x^{(\tau)}\right]^{q_x^{(i)}/q_x^{(\tau)}}$ (To get primed rates from unprimed rates from (i))

3. **Uniform Distribution of Death (UDD) for Multiple Decrement (MUDD) Table**

For any $t\in[0,1]$ and integer-valued x,

(i) ${}_tp_x^{(\tau)}\mu_{x+t}^{(i)} = q_x^{(i)}$ or equivalently $\mu_{x+t}^{(i)} = \frac{q_x^{(i)}}{1-tq_x^{(\tau)}}$for $t\neq 1$

(ii) *Ratio* Property : $\frac{{}_tq_x^{(i)}}{{}_tq_x^{(\tau)}} = \frac{\mu_{x+s}^{(i)}}{\mu_{x+s}^{(\tau)}}$ for any $s\in[0,1]$

(iii) *Partition* Property : ${}_tp_x^{\prime(i)} = \left[{}_tp_x^{(\tau)}\right]^{q_x^{(i)}/q_x^{(\tau)}}$ (To get primed rates from unprimed rates ${}_tq_x^{(i)}$ and ${}_tp_x^{(\tau)}$)

Discrete jumps: Handling Both Discrete and Continuous Decrement

1) ${}_sq_x^{(i)} = \int_0^s\left[\prod_{j=1,\,j\neq i}^{m} {}_tp_x^{\prime(j)}\right]{}_tp_x^{\prime(i)}\mu_{x+t}^{(i)}dt$ holds when decrement i is **continuous.**

2) ${}_sq_x^{(i)} = \sum_{t_k\le s}\left[\prod_{j=1,\,j\neq i}^{m} {}_{t_k}p_x^{\prime(j)}\right]\Delta({}_{t_k}q_x^{\prime(i)})$ holds when decrement i is **discrete**

Here t_k are the jump times and $\Delta\left({}_{t_k}q_x^{\prime(i)}\right)$ is the jump size at time t_k.

Case	Given	Sought	Assumption	Results
A1	$q_x'^{(1)}$ and $q_x'^{(2)}$	${}_tq_x^{(1)}$ and ${}_tq_x^{(2)}$	SUDD	$${}_tq_x^{(1)} = q_x'^{(1)}\left(t-\frac{t^2}{2}\cdot q_x'^{(2)}\right)$$ $${}_tq_x^{(2)} = q_x'^{(2)}\left(t-\frac{t^2}{2}\cdot q_x'^{(1)}\right)$$
A2	$q_x'^{(1)}$ and $q_x'^{(2)}$	$q_x^{(1)}$ and $q_x^{(2)}$	SUDD	$$q_x^{(1)} = q_x'^{(1)}\left(1-\frac{1}{2}\cdot q_x'^{(2)}\right)$$ $$q_x^{(2)} = q_x'^{(2)}\left(1-\frac{1}{2}\cdot q_x'^{(1)}\right)$$
B1	$q_x'^{(1)}$ and $q_x'^{(2)}$	${}_tq_x^{(1)}$ and ${}_tq_x^{(2)}$	CF	$${}_tq_x^{(1)} = \frac{\mu^{(1)}}{\mu^{(\tau)}}\cdot\left(1-{}_tp_x^{(\tau)}\right)$$ $${}_tq_x^{(2)} = \frac{\mu^{(2)}}{\mu^{(\tau)}}\cdot\left(1-{}_tp_x^{(\tau)}\right)$$
B2	$q_x'^{(1)}$ and $q_x'^{(2)}$	$q_x^{(1)}$ and $q_x^{(2)}$	CF	$$q_x^{(1)} = \frac{\mu^{(1)}}{\mu^{(\tau)}}\cdot q_x^{(\tau)}$$ $$q_x^{(2)} = \frac{\mu^{(2)}}{\mu^{(\tau)}}\cdot q_x^{(\tau)}$$
C1	$q_x^{(1)}$ and $q_x^{(2)}$	${}_tq_x'^{(1)}$ and ${}_tq_x'^{(2)}$	MUDD	$${}_tq_x'^{(1)} = 1-\left(1-t\cdot q_x^{(\tau)}\right)^{q_x^{(1)}/q_x^{(\tau)}}$$ $${}_tq_x'^{(2)} = 1-\left(1-t\cdot q_x^{(\tau)}\right)^{q_x^{(2)}/q_x^{(\tau)}}$$
C2	$q_x^{(1)}$ and $q_x^{(2)}$	$q_x'^{(1)}$ and $q_x'^{(2)}$	MUDD	$$q_x'^{(1)} = 1-\left(1-q_x^{(\tau)}\right)^{q_x^{(1)}/q_x^{(\tau)}}$$ $$q_x'^{(2)} = 1-\left(1-q_x^{(\tau)}\right)^{q_x^{(2)}/q_x^{(\tau)}}$$
D1	$q_x^{(1)}$ and $q_x^{(2)}$	${}_tq_x'^{(1)}$ and ${}_tq_x'^{(2)}$	CF	$${}_tq_x'^{(1)} = 1-\left({}_tp_x^{(\tau)}\right)^{{}_tq_x^{(1)}/{}_tq_x^{(\tau)}}$$ $${}_tq_x'^{(2)} = 1-\left({}_tp_x^{(\tau)}\right)^{{}_tq_x^{(2)}/{}_tq_x^{(\tau)}}$$
D2	$q_x^{(1)}$ and $q_x^{(2)}$	$q_x'^{(1)}$ and $q_x'^{(2)}$	CF	$$q_x'^{(1)} = 1-\left(p_x^{(\tau)}\right)^{q_x^{(1)}/q_x^{(\tau)}}$$ $$q_x'^{(2)} = 1-\left(p_x^{(\tau)}\right)^{q_x^{(2)}/q_x^{(\tau)}}$$
E	$q_x'^{(1)}$ and $q_x'^{(2)}$	$q_x^{(1)}$ and $q_x^{(2)}$	MUDD	$$q_x^{(1)} = q_x^{(\tau)}\cdot\left[\frac{\ln p_x'^{(1)}}{\ln p_x^{(\tau)}}\right]$$ $$q_x^{(2)} = q_x^{(\tau)}\cdot\left[\frac{\ln p_x'^{(2)}}{\ln p_x^{(\tau)}}\right]$$
F	$q_x^{(1)}$ and $q_x^{(2)}$	$q_x'^{(1)}$ and $q_x'^{(2)}$	SUDD	See Solution